# NANDROLONE DECANOATE

## PRODUCT OVERVIEW

| Specification | Details |
|---|---|
| **Generic Name** | **Nandrolone Decanoate** |
| **Drug Class** | **Anabolic Androgenic Steroid (AAS)** |
| **Active Ingredient** | **Nandrolone Decanoate** |
| **Chemical Formula** | $C_{28}H_{44}O_3$ |
| **Molecular Weight** | **428.65 g/mol** |
| **CAS Number** | **360-70-3** |
| **Administration Route** | **Intramuscular Injection** |
| **Ester Type** | **Long-Acting Nandrolone Ester** |
| **Approximate Half-Life** | **Approximately 6–12 Days** |
| **Release Profile** | **Slow and Extended Release** |
| **Bioavailability** | **High After Intramuscular Administration** |
| **Primary Medical Use** | **Treatment of selected anemia disorders, osteoporosis, and conditions associated with protein catabolism under medical supervision** |

## INTRODUCTION

Nandrolone Decanoate is a long-acting synthetic form of nandrolone developed to provide prolonged release following intramuscular administration. It is a synthetic anabolic androgenic steroid structurally related to testosterone but characterized by the presence of a decanoate ester at the 17β-hydroxyl group, which significantly increases its lipophilicity. The decanoate ester increases the lipophilicity of nandrolone, producing slower absorption from the injection site and sustained release into systemic circulation. Following administration, nandrolone decanoate is hydrolyzed by esterases to release nandrolone, which then exerts androgenic and anabolic effects through interaction with androgen receptors in target tissues. Because of its prolonged release profile compared with shorter-acting nandrolone esters, it has made Nandrolone Decanoate one of the most extensively studied anabolic steroid compounds in clinical medicine and pharmaceutical research.

## HISTORY

The discovery of nandrolone represented an important advancement in anabolic steroid research during the mid-twentieth century.

Scientists developed nandrolone by modifying the testosterone molecule through removal of the carbon atom at the 19th position, resulting in compounds with distinct anabolic and androgenic characteristics.

Subsequent esterification with decanoic acid produced Nandrolone Decanoate, allowing prolonged therapeutic activity following intramuscular administration.

Nandrolone Decanoate has been investigated for multiple medical applications involving protein metabolism, bone physiology, and hematologic abnormalities.

Scientific investigation of nandrolone contributed significantly to advances in:

- Anabolic steroid pharmacology
- Protein metabolism
- Bone remodeling
- Clinical endocrinology

## CHEMICAL STRUCTURE

Nandrolone Decanoate consists of the nandrolone molecule esterified with decanoic acid at the 17 beta hydroxyl position.

The esterification process modifies pharmacokinetic behavior without altering the biological activity of nandrolone after enzymatic hydrolysis.

The decanoate ester significantly:

- Increases lipophilicity
- Slows absorption rate
- Prolongs circulation time
- Extends duration of action

## CHEMICAL PROFILE

| Parameter | Information |
|---|---|
| **Chemical Name** | **Nandrolone Decanoate** |
| **Molecular Formula** | $C_{28}H_{44}O_3$ |
| **Molecular Weight** | **428.65 g/mol** |
| **CAS Number** | **360-70-3** |
| **Hormone Classification** | **Androgen** |
| **4-Ien Group** | **Decanoate** |
| **Active Hormone** | **Nandrolone** |

## PHARMACOLOGY

Following intramuscular administration, Nandrolone Decanoate forms a long-lasting depot within muscle tissue.

The decanoate ester undergoes gradual enzymatic hydrolysis, releasing free nandrolone into systemic circulation over an extended period.

Once released, nandrolone interacts with androgen receptors located throughout numerous tissues including:

- Skeletal muscle
- Bone
- Bone marrow
- Connective tissue
- Liver
- Kidney
- Reproductive tissues

Nandrolone also undergoes metabolic conversion into biologically active metabolites.

| Conversion Pathway | Enzyme | Product |
|---|---|---|
| **Nandrolone → DHN** | **5-alpha reductase** | **Dihydronandrolone** |
| **Nandrolone → Estradiol** | **Aromatase** | **Estradiol (Low Rate)** |

## MECHANISM OF ACTION

Nandrolone produces its biological effects primarily through activation of intracellular androgen receptors.

The biological process includes:

- Nandrolone enters target cells
- Binding to androgen receptors
- The receptor-hormone complex becomes activated
- The activated complex translocates into the cell nucleus
- Interaction with DNA regulates gene expression
- Altered protein synthesis occurs
- Cellular protein synthesis and tissue-specific physiological responses are influenced.

Activation of androgen receptors contributes to:

- Protein synthesis
- Muscle growth
- Muscle tissue maintenance
- Bone remodeling
- Red blood cell production
- Connective tissue physiology
- Endocrine regulation

## PHARMACODYNAMICS

Nandrolone Decanoate affects multiple physiological systems through androgen receptor signaling.

## SKELETAL MUSCLE SYSTEM

Nandrolone contributes to normal muscle physiology through:

- Regulation of protein synthesis
- Support of muscle tissue maintenance
- Enhancement of nitrogen retention
- Regulation of muscle cell function

Androgen receptor activation plays an important role in maintaining skeletal muscle structure and physiological function.

## SKELETAL SYSTEM

Nandrolone contributes to skeletal health by influencing:

- Bone mineral density
- Bone remodeling processes
- Osteoblast activity
- Skeletal maintenance

Nandrolone has been extensively investigated for its effects on bone metabolism and connective tissue physiology.

## HEMATOLOGICAL SYSTEM

Nandrolone influences erythropoiesis, the physiological process responsible for red blood cell production.

Physiological effects include:

- Regulation of the erythropoietic activity
- Support of red blood cell formation
- Influence on oxygen transport capacity

## CONNECTIVE TISSUE

Nandrolone contributes to connective tissue physiology by influencing:

- Collagen synthesis
- Connective tissue maintenance
- Structural protein metabolism
- Tissue remodeling

## REPRODUCTIVE SYSTEM

Nandrolone interacts with androgen-sensitive tissues and contributes to:

- Hormonal regulation
- Maintenance of androgen-responsive tissues
- Reproductive function
- Reproductive physiology

## PHARMACOKINETICS

Following intramuscular administration, Nandrolone Decanoate forms a long-lasting depot within muscle tissue.

The decanoate ester undergoes gradual enzymatic hydrolysis, releasing free nandrolone into systemic circulation over an extended period.

Compared with shorter-acting nandrolone esters, Nandrolone Decanoate provides prolonged hormone availability and sustained pharmacokinetic activity.

## PHARMACOKINETIC PROFILE

| Parameter | Description |
|---|---|
| **Ester Type** | **Long-Acting Nandrolone Ester** |
| **Absorption** | **Slow release from intramuscular depot** |
| **Release Pattern** | **Extended and controlled release** |
| **Half-Life** | **Approximately 6–12 Days** |
| **Duration of Activity** | **Long-Acting** |
| **Plasma Stability** | **Sustained nandrolone availability** |
| **Protein Binding** | **High** |
| **Metabolism** | **Primarily hepatic** |
| **Active Hormone** | **Nandrolone** |
| **Elimination** | **Urinary excretion of metabolites** |

## PHYSIOLOGICAL EFFECTS

Normal nandrolone activity contributes to:

Muscle Function
- Protein synthesis
- Muscle tissue maintenance
- Maintenance of lean body tissue
- Muscle physiology

Bone Health
- Bone remodeling
- Skeletal tissue maintenance
- Skeletal integrity

Connective Tissue
- Collagen synthesis
- Connective tissue maintenance
- Structural support

Hematological Function
- Red blood cell production
- Oxygen transport support
- Normal erythropoiesis

General Physiology
- Protein metabolism
- Nitrogen retention
- Endocrine function

## CLINICAL APPLICATIONS

Nandrolone Decanoate may be used under medical supervision for selected medical conditions when clinically indicated.

Clinical evaluation may include:

- Medical history
- Physical examination
- Laboratory assessment
- Individual therapeutic evaluation

Potential medical applications include:

- Anemia-related disorders
- Osteoporosis
- Conditions associated with protein catabolism
- Selected chronic disease management

## SAFETY CONSIDERATIONS

Nandrolone therapy requires individualized medical evaluation and regular monitoring.

Important considerations include:

- Cardiovascular health
- Liver function tests
- Endocrine function
- Liver and kidney function
- Blood pressure
- Individual therapeutic response

## POTENTIAL ADVERSE EFFECTS

Possible adverse effects may include:

Endocrine Effects
- Suppression of endogenous testosterone production
- Gynecomastia
- Alterations in endocrine function

Dermatological Effects
- Acne
- Increased oil production
- Hair loss in genetically predisposed individuals

Cardiovascular Effects
- Changes in blood lipids
- Hypertension
- Fluid retention

Reproductive Effects
- Reduced fertility
- Menstrual irregularities
- Hormonal suppression

## CONTRAINDICATIONS

| Condition | Reason |
|---|---|
| **Prostate Cancer** | **Potential androgen sensitivity** |
| **Male Breast Cancer** | **Hormone-sensitive condition** |
| **Pregnancy** | **Risk of fetal virilization** |
| **Severe Untreated Medical Conditions** | **Requires medical evaluation** |
| **Known Hypersensitivity** | **Risk of allergic reaction** |

## DRUG INTERACTIONS

Potential interactions may occur with:

Anticoagulants
Nandrolone may influence anticoagulant response and require monitoring.

Antidiabetic Medications
Changes in glucose metabolism may alter therapeutic requirements.

Corticosteroids
Combined administration may increase fluid retention.

Other Hormonal Therapies
Combined hormonal therapy requires professional medical supervision.

## LABORATORY MONITORING

| Laboratory Test | Purpose |
|---|---|
| **CBC** | **Assess blood parameters** |
| **Hematocrit** | **Monitor red blood cell concentration** |
| **Hemoglobin** | **Evaluate erythropoiesis** |
| **Liver Function Tests** | **Assess hepatic health** |
| **Kidney Function Tests** | **Evaluate renal function** |
| **Lipid Profile** | **Monitor cardiovascular risk markers** |
| **Total Testosterone** | **Evaluate endocrine status** |
| **Estradiol (E2)** | **Assess estrogen balance** |
| **PSA (When Appropriate)** | **Monitor prostate health** |

## STORAGE & STABILITY

- Store according to official product labeling.
- Protect from excessive heat and direct sunlight.
- Keep dry.
- Maintain original packaging integrity.
- Keep away from children.
- Inspect packaging before use.

## PRODUCT AUTHENTICITY

Every genuine NEXON product should include appropriate quality verification features.

Authentication may include:

- Batch number tracking
- Manufacturing information
- Expiration date verification
- Quality control documentation
- Authorized distribution information

NEXON emphasizes transparency, traceability, and product quality control.

## FREQUENTLY ASKED QUESTIONS

**Where is Nandrolone Decanoate?**
Nandrolone Decanoate is a long-acting esterified form of nandrolone designed to provide sustained hormone release following intramuscular administration.

**How is Nandrolone Decanoate different from Nandrolone Phenylpropionate?**
Nandrolone Decanoate has a longer ester chain, resulting in slower release, longer duration of activity, whereas Nandrolone Phenylpropionate produces a faster onset with a shorter duration.

**What is the active hormone released after ester hydrolysis?**
After enzymatic cleavage, the active hormone released is nandrolone (19-nortestosterone).

**Why does nandrolone activate androgen receptors?**
Nandrolone mimics the physiological effects of testosterone by binding to androgen receptors and initiating androgen-mediated cellular responses.

**Is Nandrolone Decanoate identical to testosterone?**
No. Nandrolone is a structurally modified androgen derived from testosterone by removal of the carbon atom at the 19th position, resulting in distinct pharmacological properties.

## NEXON QUALITY COMMITMENT

NEXON is committed to maintaining the highest standards of manufacturing quality, product consistency, traceability, and quality control.

Through advanced quality control systems, authentication technologies, and evidence-based pharmaceutical development, NEXON strives to provide reliable products supported by scientific transparency and regulatory standards.

## SCIENTIFIC REFERENCES


1. Bassett S. Androgen Abuse in Athletes: Detection and Consequences.

2. Kicman AT. Pharmacology of Anabolic Steroids.

3. Llewellyn W. Anabolics.

4. Kuhn CM. Anabolic Steroids.

5. Evans NA. Current Concepts in Anabolic-Androgenic Steroids.

6. Shahidi NT. A Review of the Chemistry, Biological Action and Clinical Applications of Anabolic-Androgenic Steroids.

7. Handelsman DJ. Androgen Physiology, Pharmacology and Abuse.

8. Sharan S. et al. Testosterone Therapy and Anabolic Steroid Pharmacology.

9. Nieschlag E., Behre HM. Andrology: Male Reproductive Health and Dysfunction.

10. Kicman AT, Gower DB. Anabolic Steroids in Sport and Exercise.