NEXON

TESTOSTERONE PHENYLPROPIONATE BLEND

MULTI ESTER TESTOSTERONE FORMULATION

300 mg mL

INTRAMUSCULAR INJECTION

PRODUCT OVERVIEW

Specification	Details
Generic Name	Testosterone Ester Blend
Drug Class	Androgen / Anabolic-Androgenic Steroid
Active Ingredients	Testosterone Propionate, Testosterone Phenylpropionate, Testosterone Isocaproate
Administration Route	Intramuscular Injection
Dosage Form	Oil-Based Injectable Solution
Chemical Class	Esterified Testosterone Derivatives
Primary Target	Androgen Receptor
Clinical Status	Testosterone Replacement / Androgen Therapy

INTRODUCTION

Testosterone Phenylpropionate Blend is a concentrated injectable formulation containing three esterified forms of testosterone. The selected esters possess different release characteristics, creating a combined

pharmacokinetic profile that includes both comparatively rapid and prolonged testosterone availability following intramuscular administration.

After administration, the testosterone esters are gradually released from the intramuscular depot and hydrolyzed to biologically active testosterone. The released hormone interacts with androgen receptors throughout the body and influences reproductive physiology, skeletal muscle, bone metabolism, erythropoiesis, sexual function, and numerous androgen-dependent tissues.

HISTORY

Testosterone was first isolated and characterized during the 1930s, leading to the development of esterified testosterone preparations intended to improve stability and extend biological activity after administration. Early short-acting esters were followed by longer-acting compounds that allowed testosterone to be released more gradually from an intramuscular depot.

Multi-ester testosterone preparations were subsequently developed to combine esters with different release characteristics in a single injectable formulation. These products were designed to provide an initial release of testosterone together with a more sustained phase of exposure.

Testosterone Phenylpropionate Blend represents a multi-ester formulation centered on testosterone phenylpropionate. The exact combination is not a universally standardized pharmacopoeial product, and its regulatory status depends on the manufacturer, formulation, and jurisdiction.

CHEMICAL STRUCTURE

Testosterone Phenylpropionate Blend contains three esterified derivatives of testosterone. Each component shares the same testosterone nucleus but carries a different ester group at the 17β-hydroxyl position.

The ester group affects lipid solubility, depot release, and the rate at which the compound is hydrolyzed to active testosterone.

- Three esterified testosterone components
- Shared testosterone steroid nucleus
- Oil-soluble injectable formulation
- Hydrolysis to active testosterone after release
- Combined rapid, intermediate, and prolonged release characteristics

CHEMICAL PROFILE

Parameter	Information
Primary Components	Three Testosterone Esters
Hormone Classification	Androgen / Anabolic-Androgenic Steroid
Administration	Intramuscular Injection
Dosage Form	Oil-Based Injectable Solution
Molecular Type	Esterified Steroid Hormones

Parameter	Information
Primary Receptor Target	Androgen Receptor
Testosterone Propionate	50 mg/mL
Testosterone Phenylpropionate	150 mg/mL
Testosterone Isocaproate	100 mg/mL
Total Concentration	300 mg/mL

PHARMACOLOGY

Testosterone Phenylpropionate Blend acts primarily as a source of exogenous testosterone. Following intramuscular administration, its esterified components are released from the oil depot at different rates and hydrolyzed by esterases to produce active testosterone.

Testosterone enters target cells and binds to intracellular androgen receptors. The resulting hormone-receptor complex influences gene transcription and regulates biological processes associated with male sexual development, reproductive function, skeletal muscle, bone tissue, erythropoiesis, and general androgen physiology.

Testosterone Esters → Depot Release and Ester Hydrolysis → Active Testosterone → Androgen-Receptor Activation → Androgen-Responsive Gene Expression

Testosterone may also undergo conversion to dihydrotestosterone through 5α-reductase or to estradiol through aromatase.

MECHANISM OF ACTION

The primary mechanism involves:

- Gradual release of testosterone esters from the intramuscular depot
- Enzymatic hydrolysis of each ester to active testosterone
- Binding of testosterone to intracellular androgen receptors
- Regulation of androgen-responsive gene transcription
- Conversion of testosterone to dihydrotestosterone in selected tissues
- Aromatization of testosterone to estradiol
- Modulation of protein, nitrogen, bone, and reproductive physiology

Differences between the esters primarily influence the rate and duration of testosterone release rather than the fundamental action of the released hormone.

PHARMACODYNAMICS

- Androgen-receptor signaling
- Protein metabolism

- Nitrogen retention
- Erythropoiesis
- Bone metabolism
- Sexual and reproductive physiology
- Secondary male sexual characteristics
- Endocrine feedback regulation

SKELETAL MUSCLE SYSTEM

Testosterone interacts with androgen receptors in skeletal muscle and may influence:

- Protein synthesis and metabolism
- Muscle tissue maintenance
- Nitrogen balance
- Muscle-cell differentiation
- Recovery of androgen-dependent tissue
- Maintenance of lean body mass

Androgen-receptor signaling contributes to the regulation of normal skeletal-muscle structure, function, and metabolism.

SKELETAL SYSTEM

Androgen and estrogen signaling contribute to skeletal physiology and may influence:

- Bone remodeling
- Bone mineral density
- Skeletal maintenance
- Calcium and mineral metabolism
- Structural integrity

A portion of testosterone is converted to estradiol, which also has an important role in maintaining normal bone physiology.

CONNECTIVE TISSUE

Androgen signaling may influence connective-tissue physiology through:

- Structural protein metabolism
- Tissue maintenance
- Tissue remodeling
- Collagen metabolism
- Interaction with musculoskeletal adaptation

The response of connective tissue may differ from that of skeletal muscle and depends on hormonal status, physical activity, health, and individual physiology.

REPRODUCTIVE SYSTEM

Testosterone interacts with androgen-sensitive tissues and may influence:

- Sexual function

- Libido
- Male reproductive physiology
- Accessory reproductive organs
- Secondary sexual characteristics
- Androgen-responsive tissues
- Hypothalamic-pituitary-gonadal feedback

Exogenous testosterone can suppress luteinizing hormone and follicle-stimulating hormone, potentially reducing intratesticular testosterone production and spermatogenesis.

PHARMACOKINETICS

Following intramuscular injection, the esterified testosterone components remain temporarily within an oil-based depot. Differences in ester-chain structure influence lipid solubility and the rate at which each component enters the circulation.

Testosterone propionate provides the comparatively faster-release component. Testosterone phenylpropionate forms the principal intermediate-release component, while testosterone isocaproate contributes a more sustained release phase within the selected formulation.

Once released, the esters are hydrolyzed to active testosterone. Circulating testosterone is extensively bound to sex hormone-binding globulin and albumin. Metabolism occurs mainly in the liver and peripheral tissues, producing active metabolites such as dihydrotestosterone and estradiol as well as inactive metabolites eliminated primarily through urine.

The complete pharmacokinetic profile of this specific 300 mg/mL combination has not been established through standardized clinical studies.

PHYSIOLOGICAL EFFECTS

Testosterone plays a central role in male reproductive development, sexual function, maintenance of secondary sexual characteristics, skeletal-muscle physiology, bone metabolism, erythropoiesis, and overall androgen-dependent health.

When testosterone concentrations are medically restored in individuals with confirmed androgen deficiency, physiological effects may include maintenance of sexual function, lean tissue, bone density, mood, energy, and red-blood-cell production. The magnitude and clinical relevance of these effects depend on the underlying diagnosis and individual response.

Exposure above normal physiological concentrations can increase the likelihood of endocrine suppression, erythrocytosis, altered lipid parameters, fluid retention, acne, androgen-related hair loss, fertility impairment, and cardiovascular complications. Effects vary according to concentration, duration, formulation, and patient characteristics.

CLINICAL APPLICATIONS

Approved testosterone preparations are used principally as replacement therapy in appropriately evaluated patients with confirmed testosterone deficiency associated with specific medical conditions.

Clinical assessment generally requires compatible symptoms together with consistently low serum testosterone concentrations measured according to established medical guidelines. Treatment selection depends on diagnosis, age, comorbidities, fertility considerations, formulation characteristics, and monitoring requirements.

This specific three-ester 300 mg/mL formulation is not a universally standardized or independently FDA-approved medicine. Its clinical and regulatory status must therefore be distinguished from that of individually approved testosterone products.

SAFETY CONSIDERATIONS

Testosterone exposure requires consideration of cardiovascular, hematological, endocrine, reproductive, metabolic, prostate, hepatic, and psychological health.

Important safety considerations include:

- Cardiovascular health
- Blood pressure
- Hematocrit and hemoglobin
- Lipid profile
- Liver function
- Endocrine function
- Reproductive function and fertility
- Prostate health
- Sleep apnea risk
- Fluid retention
- Injection-site reactions

The overall safety profile depends on the formulation, concentration, duration of exposure, individual response, and patient characteristics. Medical evaluation and regular monitoring are important whenever testosterone therapy is clinically indicated.

POTENTIAL ADVERSE EFFECTS

Potential adverse effects associated with testosterone and other anabolic-androgenic steroids may involve endocrine, cardiovascular, hematological, dermatological, reproductive, hepatic, metabolic, and psychological systems.

Endocrine Effects

- Suppression of endogenous testosterone production
- Suppression of luteinizing hormone and follicle-stimulating hormone
- Alterations in estrogen and androgen balance
- Gynecomastia in susceptible individuals
- Testicular atrophy
- Potential suppression of spermatogenesis

Cardiovascular Effects

- Changes in lipid profile
- Possible increases in blood pressure
- Fluid and sodium retention
- Potential thromboembolic complications
- Possible increase in cardiovascular risk

Hematological Effects

- Increased hematocrit
- Increased hemoglobin
- Erythrocytosis or polycythemia
- Increased blood viscosity

Dermatological Effects

- Acne
- Increased sebaceous activity
- Oily skin
- Androgen-related hair loss in genetically predisposed individuals
- Increased body or facial hair

Reproductive Effects

- Reduced sperm production
- Changes in fertility
- Alterations in reproductive hormone regulation
- Changes in libido or sexual function
- Prostate-related effects

Hepatic and Metabolic Effects

- Changes in lipid parameters
- Alterations in glucose metabolism
- Changes in liver-function parameters
- Fluid and electrolyte imbalance
- Changes in body composition

Administration Related Effects

- Injection-site pain
- Swelling or irritation
- Inflammatory reactions
- Hypersensitivity to active ingredients or excipients
- Infection when administration is not performed under appropriate sterile conditions

The occurrence and severity of adverse effects vary according to exposure, concentration, duration, individual response, underlying health conditions, and concurrent medications.

CONTRAINDICATIONS

Condition	Reason
Known or Suspected Prostate Carcinoma	Androgen-sensitive disease may be stimulated
Known or Suspected Male Breast Carcinoma	Potential androgen-sensitive malignancy
Pregnancy	Testosterone may cause fetal harm

Condition	Reason
Breastfeeding	Insufficient safety and potential hormonal exposure
Known Hypersensitivity	Risk of allergic reaction to testosterone esters or excipients
Unexplained Elevated Hematocrit	Increased risk of erythrocytosis-related complications
Severe Uncontrolled Cardiovascular Disease	Potential fluid, blood-pressure, and cardiovascular effects
Severe Hepatic or Renal Disease	Potential fluid retention and impaired clinical tolerance

DRUG INTERACTIONS

Potential interactions or important clinical considerations may exist with:

- Anticoagulants
- Insulin and antidiabetic medications
- Corticosteroids
- Adrenocorticotropic hormone
- Other androgenic or anabolic agents
- Medications affecting blood pressure
- Drugs influencing fluid and electrolyte balance
- Hormonal therapies
- Laboratory tests affected by androgen status

Testosterone may alter anticoagulant response, glucose-control requirements, fluid retention, and selected laboratory measurements. The clinical significance depends on the specific medication, health condition, formulation, exposure, and individual response.

LABORATORY MONITORING

Laboratory Test	Purpose
CBC	Monitor hematological parameters
Hematocrit / Hemoglobin	Assess erythrocytosis
ALT / AST	Liver-function assessment
Lipid Profile	Cardiovascular-risk assessment
Blood Pressure	Cardiovascular monitoring
Total Testosterone	Assess testosterone exposure
Free Testosterone / SHBG	Evaluate bioavailable androgen status
Estradiol	Assess estrogenic conversion when clinically indicated

Laboratory Test	Purpose
LH / FSH	Assess hypothalamic-pituitary-gonadal suppression
PSA	Prostate monitoring when clinically appropriate
Fasting Glucose / HbA1c	Metabolic assessment

STORAGE AND STABILITY

- Store according to official product labeling.
- Protect from excessive heat and direct sunlight.
- Do not freeze.
- Maintain original packaging integrity.
- Keep away from children.
- Inspect the solution and packaging before use.
- Do not use if the container is damaged or the solution shows unexpected contamination.
- Crystallization and storage requirements depend on the validated formulation.

PRODUCT AUTHENTICITY

Every genuine NEXON product should include appropriate quality-verification features.

- Batch-number tracking
- Manufacturing information
- Expiration-date verification
- QR authentication system
- Tamper-evident packaging
- Authorized-distribution verification

NEXON emphasizes transparency, traceability, and product-quality control.

FREQUENTLY ASKED QUESTIONS

What is Testosterone Phenylpropionate Blend?

Testosterone Phenylpropionate Blend is a concentrated oil-based injectable formulation containing three testosterone esters with different release characteristics.

Which testosterone esters are included?

The selected formulation contains testosterone propionate, testosterone phenylpropionate, and testosterone isocaproate, with phenylpropionate serving as the primary component.

What does 300 mg/mL represent?

It represents the combined labeled concentration of the three esterified testosterone ingredients: 300 mg per mL.

Is Testosterone Phenylpropionate Blend a peptide?

No. It is an esterified androgen formulation and belongs to the anabolic-androgenic steroid class.

Is Testosterone Phenylpropionate Blend a SARM?

No. Testosterone is a steroid hormone that activates androgen receptors and is not classified as a selective androgen-receptor modulator.

Is Testosterone Phenylpropionate Blend oral or injectable?

It is presented as an oil-based solution intended for intramuscular administration.

Why are multiple testosterone esters combined?

Different ester structures produce different depot-release characteristics. Combining them creates a formulation with comparatively rapid, intermediate, and prolonged release phases.

Do the individual esters have different biological actions?

After hydrolysis, all three esters release active testosterone. Their principal difference is the rate and duration of release rather than the fundamental action of testosterone.

Can Testosterone Phenylpropionate Blend suppress natural testosterone production?

Yes. Exogenous testosterone can suppress hypothalamic-pituitary-gonadal signaling, including luteinizing hormone, follicle-stimulating hormone, endogenous testosterone production, and spermatogenesis.

Is this specific blend an FDA-approved formulation?

No independent FDA approval has been established for this specific three-ester, 300 mg/mL combination. Regulatory status varies according to the actual manufacturer and jurisdiction.

NEXON QUALITY COMMITMENT

NEXON is committed to maintaining high standards of manufacturing quality, product consistency, traceability, and scientific transparency.

Through quality-control systems, authentication procedures, and responsible information sharing, NEXON aims to provide reliable pharmaceutical products and educational resources for healthcare professionals and informed users.

SCIENTIFIC REFERENCES

- Nieschlag E, Behre HM, Nieschlag S. Testosterone: Action, Deficiency, Substitution.
- Nieschlag E, Behre HM. Andrology: Male Reproductive Health and Dysfunction.
- Bhasin S, et al. Testosterone Therapy in Men With Hypogonadism: An Endocrine Society Clinical Practice Guideline.
- Kicman AT. Pharmacology of Anabolic Steroids.
- Handelsman DJ. Androgen Physiology, Pharmacology, Use and Misuse.
- Basaria S. Androgen Abuse in Athletes: Detection and Consequences.
- Shahidi NT. A Review of the Chemistry, Biological Action, and Clinical Applications of Anabolic-Androgenic Steroids.
- Mooradian AD, Morley JE, Korenman SG. Biological Actions of Androgens.

- Snyder PJ. Clinical Use of Androgens.
- Wilson JD. Androgen Physiology and Pharmacology.
- Swerdloff RS, Wang C. Androgen Deficiency and Testosterone Replacement.
- U.S. Food and Drug Administration. Prescribing Information for Approved Injectable Testosterone Products.