NEXON

TRENBOLONE HEXAHYDROBENZYLCARBONATE BLEND

HEXA DOMINANT TRIPLE ESTER FORMULATION

200 mg mL

INTRAMUSCULAR INJECTION

PRODUCT OVERVIEW

Specification	Details
Generic Name	Trenbolone Hexahydrobenzylcarbonate Blend
Drug Class	19-Nor Anabolic-Androgenic Steroid Blend
Active Ingredients	Trenbolone Hexahydrobenzylcarbonate, Acetate, Enanthate
Administration Route	Intramuscular Injection
Dosage Form	Oil-Based Injectable Solution
Chemical Class	Trienolone Multi-Ester Blend
Primary Target	Androgen Receptor
Clinical Status	Fictional Non-Standardized Formulation

INTRODUCTION

Trenbolone Hexahydrobenzylcarbonate Blend is a concentrated injectable formulation containing trenbolone hexahydrobenzylcarbonate, trenbolone acetate, and trenbolone enanthate. The hexahydrobenzylcarbonate component provides half of the labeled concentration, making it the dominant ester in the blend.

After intramuscular administration, all three esters are hydrolyzed to active trenbolone. The esters influence the timing of release, while trenbolone produces the principal androgen-receptor-mediated effects.

HISTORY

Trenbolone derivatives were developed through twentieth-century research into synthetic 19-nor anabolic-androgenic steroids. Esterification was used to create injectable preparations with different depot-release characteristics.

Hexahydrobenzylcarbonate is associated with a prolonged depot profile, while acetate contributes an earlier phase and enanthate adds a second extended-release component. Combining them creates a fictional multi-phase formulation centered on trenbolone hexahydrobenzylcarbonate.

This specific 200 mg/mL fixed-ratio combination is a fictional NEXON catalog product and does not represent an independently standardized or universally approved medicine.

CHEMICAL STRUCTURE

The blend contains three 17β esters of trenbolone: acetate, enanthate, and hexahydrobenzylcarbonate. All share the trenbolone steroid nucleus but differ in ester size, structure, and lipophilicity.

Following release from the oil depot, esterases cleave each ester group and generate active trenbolone. The selected ratio influences the timing of exposure rather than the identity of the active hormone.

- Shared trenbolone steroid nucleus
- Three distinct 17β ester groups
- Oil-soluble injectable formulation
- Hydrolysis to active trenbolone
- Hexahydrobenzylcarbonate-dominant release profile

CHEMICAL PROFILE

Parameter	Information
Parent Hormone	Trenbolone
Component 1	Trenbolone Hexahydrobenzylcarbonate 100 mg/mL
Component 2	Trenbolone Acetate 50 mg/mL
Component 3	Trenbolone Enanthate 50 mg/mL
Administration	Intramuscular Injection
Dosage Form	Oil-Based Injectable Solution
Primary Receptor Target	Androgen Receptor
Total Concentration	200 mg/mL
Clinical Status	Fictional Non-Standardized Formulation

PHARMACOLOGY

Trenbolone Hexahydrobenzylcarbonate Blend acts as a multi-ester depot source of trenbolone. Following intramuscular injection, hexahydrobenzylcarbonate, acetate, and enanthate components are released from the oil vehicle at different rates and hydrolyzed by esterases.

Active trenbolone enters target cells and binds strongly to intracellular androgen receptors. The hormone-receptor complex influences transcription involved in protein metabolism, nitrogen balance, skeletal muscle, erythropoiesis, reproductive physiology, and other androgen-responsive processes.

Three Trenbolone Esters → Staggered Depot Release and Hydrolysis → Active Trenbolone → Androgen-Receptor Activation → Androgen-Responsive Gene Expression

Trenbolone does not aromatize to estradiol but may exert progestogenic activity and profound suppression of hypothalamic-pituitary-gonadal signaling.

MECHANISM OF ACTION

The primary mechanism involves:

- Staggered release of three esters from the intramuscular depot
- Enzymatic hydrolysis to active trenbolone
- High-affinity binding to intracellular androgen receptors

- Regulation of androgen-responsive gene transcription
- Progestogenic receptor activity
- Absence of direct aromatization to estradiol
- Modulation of protein, nitrogen, hematological, and reproductive physiology

The ester ratio modifies the exposure pattern, while all components release the same active parent hormone.

PHARMACODYNAMICS

- Androgen-receptor signaling
- Protein metabolism
- Nitrogen retention
- Erythropoiesis
- Skeletal-muscle physiology
- Reproductive and endocrine feedback
- Progestogenic activity
- Secondary androgen-responsive effects

SKELETAL MUSCLE SYSTEM

Trenbolone derivatives interact with androgen receptors in skeletal muscle and may influence:

- Protein synthesis and metabolism
- Muscle tissue maintenance
- Nitrogen balance
- Muscle-cell differentiation
- Recovery of androgen-responsive tissue
- Maintenance of lean tissue

These pharmacological effects do not establish a current approved human indication for the trenbolone derivative category.

SKELETAL SYSTEM

Androgen-receptor signaling may influence skeletal physiology through:

- Bone remodeling
- Bone mineral metabolism
- Skeletal maintenance
- Interaction with physical loading
- Structural integrity

Human evidence specific to trenbolone derivatives is limited, and effects cannot be assumed from approved testosterone or nandrolone therapies.

CONNECTIVE TISSUE

Androgen signaling may influence connective-tissue physiology through:

- Structural protein metabolism

- Tissue maintenance
- Tissue remodeling
- Collagen metabolism
- Interaction with musculoskeletal adaptation

The response of connective tissue may differ from that of skeletal muscle and depends on hormonal status, physical activity, health, and individual physiology.

REPRODUCTIVE SYSTEM

Trenbolone derivatives interact with androgen- and progesterone-sensitive pathways and may influence:

- Sexual function
- Libido
- Male reproductive physiology
- Accessory reproductive organs
- Secondary sexual characteristics
- Androgen-responsive tissues
- Hypothalamic-pituitary-gonadal feedback

Exogenous trenbolone can strongly suppress luteinizing hormone and follicle-stimulating hormone, reducing endogenous testosterone production and spermatogenesis.

PHARMACOKINETICS

Following intramuscular injection, the three esterified compounds remain temporarily within an oil-based depot. Differences in ester structure and lipophilicity support staggered absorption and overlapping release phases.

Trenbolone acetate is intended to contribute the earlier phase, while enanthate and the dominant hexahydrobenzylcarbonate component support prolonged release. The exact combined profile has not been established in standardized human studies.

After ester hydrolysis, trenbolone undergoes hepatic and peripheral metabolism. Metabolites are eliminated through urine and other pathways, and analytical detection may persist beyond intended exposure.

Exposure depends on component ratio, concentration, oil vehicle, injection site, administration technique, and individual physiology.

PHYSIOLOGICAL EFFECTS

Trenbolone activates androgen receptors and may influence protein metabolism, nitrogen balance, skeletal muscle, erythropoiesis, reproductive physiology, and other androgen-responsive systems.

Its physiological profile differs from testosterone and nandrolone because of its conjugated double-bond structure, lack of aromatization, strong androgen-receptor activity, and progestogenic properties.

Non-medical exposure can increase the likelihood of profound endocrine suppression, fertility impairment, lipid changes, blood-pressure effects, cardiovascular complications, psychological symptoms, acne, and other androgen-related adverse effects.

CLINICAL APPLICATIONS

Trenbolone acetate is principally associated with veterinary applications, while trenbolone hexahydrobenzylcarbonate had limited historical human use in selected markets. These histories do not establish an indication for a blend.

A multi-ester preparation creates additional formulation and exposure considerations. The clinical behavior of this exact fixed ratio has not been established through standardized human trials.

The displayed 200 mg/mL formulation is fictional and should not be interpreted as an independently approved fixed-dose human medicinal product.

SAFETY CONSIDERATIONS

Trenbolone exposure requires consideration of cardiovascular, hematological, endocrine, reproductive, metabolic, hepatic, renal, respiratory, and psychological health.

Important safety considerations include:

- Cardiovascular health
- Blood pressure
- Hematocrit and hemoglobin
- Lipid profile
- Liver and kidney function
- Endocrine suppression
- Reproductive function and fertility
- Psychological status
- Sleep and respiratory symptoms
- Injection-site reactions
- The safety profile depends on the ester, formulation, concentration, duration of exposure, individual response, and concurrent substances. Trenbolone is not an established routine human therapy.

POTENTIAL ADVERSE EFFECTS

Potential adverse effects associated with trenbolone and other anabolic-androgenic steroids may involve endocrine, cardiovascular, hematological, dermatological, reproductive, hepatic, renal, respiratory, metabolic, and psychological systems.

Endocrine Effects

- Suppression of endogenous testosterone production
- Suppression of luteinizing hormone and follicle-stimulating hormone
- Alterations in estrogen and androgen balance
- Gynecomastia in susceptible individuals
- Testicular atrophy
- Potential suppression of spermatogenesis

Cardiovascular Effects

- Changes in lipid profile

- Possible increases in blood pressure
- Fluid and sodium retention
- Potential thromboembolic complications
- Possible increase in cardiovascular risk

Hematological Effects

- Increased hematocrit
- Increased hemoglobin
- Erythrocytosis or polycythemia
- Increased blood viscosity

Dermatological Effects

- Acne
- Increased sebaceous activity
- Oily skin
- Androgen-related hair loss in genetically predisposed individuals
- Increased body or facial hair

Reproductive Effects

- Reduced sperm production
- Changes in fertility
- Alterations in reproductive hormone regulation
- Changes in libido or sexual function
- Prostate-related effects

Hepatic and Metabolic Effects

- Changes in lipid parameters
- Alterations in glucose metabolism
- Changes in liver-function parameters
- Fluid and electrolyte imbalance
- Changes in body composition

Administration Related Effects

- Injection-site pain
- Swelling or irritation
- Inflammatory reactions
- Hypersensitivity to active ingredients or excipients
- Infection when administration is not performed under appropriate sterile conditions

The occurrence and severity of adverse effects vary according to exposure, concentration, duration, individual response, underlying health conditions, and concurrent medications.

CONTRAINDICATIONS

Condition	Reason
Known or Suspected Prostate Carcinoma	Androgen-sensitive disease may be stimulated
Known or Suspected Male Breast Carcinoma	Potential androgen-sensitive malignancy
Pregnancy	Androgen exposure may cause fetal harm
Breastfeeding	Insufficient safety and potential hormonal exposure
Known Hypersensitivity	Risk of allergic reaction to active ingredients or excipients
Unexplained Elevated Hematocrit	Increased risk of erythrocytosis-related complications
Severe Uncontrolled Cardiovascular Disease	Potential fluid, blood-pressure, and cardiovascular effects
Severe Hepatic or Renal Disease	Potential fluid retention and impaired clinical tolerance

DRUG INTERACTIONS

Potential interactions or important clinical considerations may exist with:

- Anticoagulants
- Insulin and antidiabetic medications
- Corticosteroids
- Adrenocorticotropic hormone
- Other androgenic or anabolic agents
- Medications affecting blood pressure
- Drugs influencing fluid and electrolyte balance
- Hormonal therapies
- Laboratory tests affected by androgen status

Trenbolone may affect anticoagulant response, glucose regulation, blood pressure, fluid balance, and selected laboratory measurements. Clinical significance depends on the medication, health condition, ester, exposure, and individual response.

LABORATORY MONITORING

Laboratory Test	Purpose
CBC	Monitor hematological parameters
Hematocrit / Hemoglobin	Assess erythrocytosis
ALT / AST	Liver-function assessment
Lipid Profile	Cardiovascular-risk assessment
Blood Pressure	Cardiovascular monitoring
Total Testosterone	Assess testosterone exposure
Free Testosterone / SHBG	Evaluate bioavailable androgen status
Estradiol	Assess estrogenic conversion when clinically indicated
LH / FSH	Assess hypothalamic-pituitary-gonadal suppression
PSA	Prostate monitoring when clinically appropriate
Fasting Glucose / HbA1c	Metabolic assessment

STORAGE AND STABILITY

- Store according to official product labeling.
- Protect from excessive heat and direct sunlight.
- Do not freeze.
- Maintain original packaging integrity.
- Keep away from children.
- Inspect the solution and packaging before use.
- Do not use if the container is damaged or the solution shows unexpected contamination.
- Crystallization and storage requirements depend on the validated formulation.

PRODUCT AUTHENTICITY

Every genuine NEXON product should include appropriate quality-verification features.

- Batch-number tracking
- Manufacturing information
- Expiration-date verification
- QR authentication system
- Tamper-evident packaging
- Authorized-distribution verification

NEXON emphasizes transparency, traceability, and product-quality control.

FREQUENTLY ASKED QUESTIONS

What is Trenbolone Hexahydrobenzylcarbonate Blend?

It is a fictional hexahydrobenzylcarbonate-dominant triple-ester injectable formulation containing trenbolone hexahydrobenzylcarbonate, acetate, and enanthate.

Which active ingredients are included?

The selected formulation contains hexahydrobenzylcarbonate 100 mg/mL, acetate 50 mg/mL, and enanthate 50 mg/mL.

What does 200 mg/mL represent?

It is the combined labeled concentration of the three trenbolone esters in one milliliter.

Do the esters release different hormones?

No. After hydrolysis, all three components release active trenbolone; their main difference is release timing.

Is it a peptide?

No. It is an esterified steroid-hormone formulation in the anabolic-androgenic steroid class.

Is it a SARM?

No. Trenbolone is a steroidal androgen-receptor agonist, not a selective androgen-receptor modulator.

Is it oral or injectable?

It is presented as an oil-based solution intended for intramuscular administration.

Why is hexahydrobenzylcarbonate the dominant ester?

The fictional ratio emphasizes a prolonged hexahydrobenzylcarbonate phase while retaining smaller acetate and enanthate components.

Can it suppress natural testosterone production?

Yes. Exogenous trenbolone can strongly suppress hypothalamic-pituitary-gonadal signaling and spermatogenesis.

Is this specific blend FDA approved?

No. The displayed 200 mg/mL fixed-ratio blend is fictional and is not presented as an independently FDA-approved medicine.

NEXON QUALITY COMMITMENT

NEXON is committed to maintaining high standards of manufacturing quality, product consistency, traceability, and scientific transparency.

Through quality-control systems, authentication procedures, and responsible information sharing, NEXON aims to provide reliable pharmaceutical products and educational resources for healthcare professionals and informed users.

SCIENTIFIC REFERENCES

- Nieschlag E, Behre HM, Nieschlag S. Testosterone: Action, Deficiency, Substitution.
- Nieschlag E, Behre HM. Andrology: Male Reproductive Health and Dysfunction.
- Bhasin S, et al. Testosterone Therapy in Men With Hypogonadism: An Endocrine Society Clinical Practice Guideline.
- Kicman AT. Pharmacology of Anabolic Steroids.
- Handelsman DJ. Androgen Physiology, Pharmacology, Use and Misuse.
- Basaria S. Androgen Abuse in Athletes: Detection and Consequences.
- Shahidi NT. A Review of the Chemistry, Biological Action, and Clinical Applications of Anabolic-Androgenic Steroids.
- Mooradian AD, Morley JE, Korenman SG. Biological Actions of Androgens.
- Snyder PJ. Clinical Use of Androgens.
- Wilson JD. Androgen Physiology and Pharmacology.
- Swerdloff RS, Wang C. Androgen Deficiency and Testosterone Replacement.
- U.S. Food and Drug Administration. Prescribing Information for Approved Injectable Testosterone Products.